



Programme Area: Bioenergy

Project: Characterisation of Feedstocks

Title: D6 Final Report (Phase 1) Appendix 8 Part 1

Abstract:

The primary objective of this 2015/16/17 Project was to provide an understanding of UK produced biomass properties, how these vary and what causes this variability.

This document is one of the appendices to the Final Report from the first Phase (2015/16) of the Characterisation of Feedstocks (CofF) project, Deliverable D6. D6 is provided in a number of parts consisting of the main body text plus 13 Appendices, provided in 17 files. These 13 appendices are provided in 12 pdf files plus 46 data files in Microsoft Excel format. The purpose of this report plus its related parts is to report the variability in feedstock properties of UK produced energy biomass, the causes of these variations and the relationship between the feedstock properties and the provenance data collected. Five feedstocks were studied: Miscanthus, willow short rotation coppice (SRC), poplar SRC, poplar grown as short rotation forests (SRF), and spruce SRF, with poplar and Sitka spruce selected to represent broadleaved and coniferous biomass crops respectively. Provenance data include site properties (such as general climate zone and soil chemistry), the conditions at the time of sample collection, and past management of the site and crop with soil samples also collected for analysis. The feedstock samples were analysed in UKAS accredited laboratories.

Context:


The Characterisation of Feedstocks project provides an understanding of UK produced 2nd generation energy biomass properties, how these vary and what causes this variability. In this project, several types of UK-grown biomass, produced under varying conditions, were sampled. The biomass sampled included Miscanthus, Short Rotation Forestry (SRF) and Short Rotation Coppice (SRC) Willow. The samples were tested to an agreed schedule in an accredited laboratory. The results were analysed against the planting, growing, harvesting and storage conditions (i.e. the provenance) to understand what impacts different production and storage methods have on the biomass properties. The main outcome of this project is a better understanding of the key characteristics of UK biomass feedstocks (focusing on second generation) relevant in downstream energy conversion applications, and how these characteristics vary by provenance.

Disclaimer:


The Energy Technologies Institute is making this document available to use under the Energy Technologies Institute Open Licence for Materials. Please refer to the Energy Technologies Institute website for the terms and conditions of this licence. The Information is licensed 'as is' and the Energy Technologies Institute excludes all representations, warranties, obligations and liabilities in relation to the Information to the maximum extent permitted by law. The Energy Technologies Institute is not liable for any errors or omissions in the Information and shall not be liable for any loss, injury or damage of any kind caused by its use. This exclusion of liability includes, but is not limited to, any direct, indirect, special, incidental, consequential, punitive, or exemplary damages in each case such as loss of revenue, data, anticipated profits, and lost business. The Energy Technologies Institute does not guarantee the continued supply of the Information. Notwithstanding any statement to the contrary contained on the face of this document, the Energy Technologies Institute confirms that the authors of the document have consented to its publication by the Energy Technologies Institute.

Appendix 8 part 1: *Miscanthus* Study 1 graphs


Moisture Content of *Miscanthus*




Net Calorific Value of *Miscanthus*




Gross Calorific Value (DAF) of *Miscanthus*




Dry Ash content of *Miscanthus*




Dry nitrogen content of *Miscanthus*




Dry chlorine content of *Miscanthus*




Barium content of *Miscanthus*




Beryllium content of *Miscanthus*




Chromium content of *Miscanthus*




Cobalt content of *Miscanthus*




Copper content of *Miscanthus*




Molybdenum content of *Miscanthus*




Nickel content of *Miscanthus*




Vanadium content of *Miscanthus*




Zinc content of *Miscanthus*




Arsenic content of *Miscanthus*




Antimony content of *Miscanthus*




Mercury content of *Miscanthus*




Cadmium content of *Miscanthus*




Lead content of *Miscanthus*




Fluorine content of *Miscanthus*




Bromine content of *Miscanthus*




Variation in ash composition of *Miscanthus*




Aluminium content of *Miscanthus*




Calcium content of *Miscanthus*




Iron content of *Miscanthus*




Potassium content of *Miscanthus*




Magnesium content of *Miscanthus*




Manganese content of *Miscanthus*




Sodium content of *Miscanthus*



Phosphorous content of *Miscanthus*



Silicon content of *Miscanthus*



Titanium content of *Miscanthus*

